



ENVISAT ASAR satellite offshore wind resource statistics in Iceland compared to NORA10 model results

Hasager, Charlotte Bay; Badger, Merete; Rugaard Furevik, Birgitte; Nawri, Nikolai; Petersen, Guðrún
Nína ; Björnsson, Halldór ; Bingöl, Ferhat; Clausen, Niels-Erik

Published in:
Geophysical Research Abstracts

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Hasager, C. B., Badger, M., Rugaard Furevik, B., Nawri, N., Petersen, G. N., Björnsson, H., Bingöl, F., & Clausen, N-E. (2014). ENVISAT ASAR satellite offshore wind resource statistics in Iceland compared to NORA10 model results. *Geophysical Research Abstracts*, 16, [EGU2014-1230].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



ENVISAT ASAR satellite offshore wind resource statistics in Iceland compared to NORA10 model results

Charlotte Bay Hasager (1), Merete Badger (1), Birgitte Rugaard Furevik (2), Nikolai Nawri (3), Guðrún Nína Petersen (3), Halldór Björnsson (3), Ferhat Ferhat Bingol (1), and Niels-Erik Clausen (1)

(1) DTU Wind Energy, Roskilde, Denmark (cbha@dtu.dk), (2) Norwegian Meteorological Institute, Bergen, Norway (birgitte.furevik@met.no), (3) Icelandic Meteorological Office, Reykjavik, Iceland (gnp@vedur.is)

In Iceland the offshore regions have been investigated from satellite images from the Envisat Advanced Synthetic Aperture Radar (SAR) from the European Space Agency. The data have been retrieved from 2002 to 2012. Each satellite image has been calibrated and thereafter the CMOD5.N geophysical model function has been used to calculate the wind speed. Each resulting pixel is 1 km by 1 km. The wind direction is taken from the US Navy Operational Global Atmospheric Prediction System (NOGAPS) model. It has much lower spatial resolution and therefore the model wind directions are interpolated in space and time before performing the SAR-based wind retrieval. In total 2,500 images have been processed. This results in average 300 overlapping images. More data were available in the northern part of Iceland; therefore the number of overlapping samples is around 400 here and only 200 at the southern coast. The wind resource statistics of mean wind speed, Weibull scale and shape parameters and energy density have been calculated using the Satellite-WAsP (S-WAsP) program. The coastline of Iceland is complex. The individual wind maps from SAR reveal a multitude of atmospheric phenomena including lee effects and gap flows in the fjords. The wind resource statistics shows the mean wind speed to range from 5 to 8 m/s at 10 m height above the sea level. Selected case study areas are being defined for further investigation. SAR-derived wind maps have the advantage of covering the coastal zone. Further offshore the SAR-derived winds will be compared to the NORA10 atmospheric model results and scatterometer winds. In Iceland the wind resources on land are promising for wind energy application but it is not yet exploited. The study on the offshore wind resource is useful as pre-feasibility in case this clean energy resource is to be exploited at a later stage. The work is part of the Nordic Icewind project.